

Orgamox™ Amoxicillin

PRESENTATION:

Orgamox™ 250 Capsule: Each capsule contains Amoxicillin Trihydrate BP equivalent to Amoxicillin 250 mg.

Orgamox™ 500 Capsule: Each capsule contains Amoxicillin Trihydrate BP equivalent to Amoxicillin 500 mg.

DESCRIPTION:

Amoxicillin is a broad spectrum bactericidal antibiotic active against both Gram-Positive and Gram-Negative bacteria.

INDICATION:

Orgamox is indicated for the treatment of following infections when caused by susceptible organisms.

RESPIRATORY TRACT INFECTIONS: Acute and chronic bronchitis, pneumonia, otitis media, sinusitis, tonsillitis, pharyngitis and laryngitis, lobar and bronchopneumonia, chronic bronchial sepsis.

SKIN AND SOFT TISSUE INFECTIONS: Cellulitis, infected wounds and abscesses, furunculosis, carbuncles, Orgamox™ may also be used as prophylactic cover for patients at risk of developing endocarditis when undergoing dental surgery.

GENITO-URINARY TRACT INFECTIONS: Pyelonephritis, cystitis, nephritis, pyelitis, prostatitis, urethritis and gonorrhoea.

GASTRO-INTESTINAL TRACT INFECTIONS: Bacillary dysentery, enteritis, peritonitis, typhoid and paratyphoid.

VENEREAL INFECTIONS: Gonorrhoea and syphilis.

OTHER INFECTIONS: Septicaemia, endocarditis, osteomyelitis, meningitis, septic abortion, intra-abdominal sepsis, dental infections and post operative use may reduce the incidence of certain post operative infections.

DOSAGE AND ADMINISTRATION:

Adult: Usual Oral Dose: 250 mg three times daily, increasing to 500 mg three times daily for more severe infections.

Children: Oral usual dose 125 mg three times daily for up to 10 years age, increasing to 250 mg three times daily for more severe infections.

SIDE-EFFECTS:

Side effects as with other penicillins, are uncommon and mainly of a mild and transitory nature; they may include diarrhoea, indigestion or occasionally rash, either urticarial or erythematous.

CONTRAINDICATION

It is contraindicated for patients hypersensitive to penicillin.

PRECAUTION

In severe renal impairment total daily dose should be reduced by 25-50%

USE IN PREGNANCY & LACTATION

Animal studies with Amoxicillin have shown no teratogenic effects. Its suitability in human pregnancy has been well documented in clinical studies. When antibiotic therapy is required during pregnancy, Amoxicillin may be considered appropriately. During lactation, trace quantities of Amoxicillin can be detected in breast milk.

DRUG INTERACTION

Concurrent administration of probenecid delays the excretion of Amoxicillin.

PHARMACEUTICAL PRECAUTION

Store in a cool dry place and protected from light. Keep all medicines out of reach of children.

PACKAGING:

Orgamox™ 250 Capsule: Box containing 100 capsules in blister pack.

Orgamox™ 500 Capsule: Box containing 50 capsules in blister pack.

TM Registered Trade Mark



Manufactured By:
Apex Pharma Ltd. for
Organic Health Care Ltd.
Gazipur, Bangladesh.